Food and Drug Administration, HHS

too slow. Animals which cannot drink after initial intravenous therapy may need to be dosed with a stomach tube or esophageal tube. Adequate colostrum intake during the first 12 hours is essential for healthy, vigorous calves. Antibacterial therapy is often indicated in bacterial scours due to E. coli and/or Salmonella. The product does not contain antibacterial agents. A veterinarian should be consulted in severely scouring calves or cases requiring antibacterial therapy. The product is not nutritionally complete if administered by itself for long periods of time. It should not be administered beyond the recommended treatment period without the addition of milk or milk replacer.

[48 FR 38606, Aug. 25, 1983, as amended at 56 FR 50653, Oct. 8, 1991; 60 FR 55659, Nov. 2, 1995]

§ 520.563 Diatrizoate meglumine and diatrizoate sodium oral solution.

- (a) Specifications. Diatrizoate meglumine oral solution is a water soluble radiopaque medium containing 66 percent diatrizoate meglumine and 10 percent diatrizoate sodium.
- (b) *Sponsor*. See No. 053501 in §510.600(c) of this chapter.
- (c) Conditions of use. (1) It is indicated for radiography of the gastrointestinal tract in dogs and cats.
- (2) It is administered orally at a dosage level of 0.5 to 1.0 milliliter per pound of body weight by gavage or stomach tube. It is administered rectally at a dosage level of 0.5 to 1.0 milliliter per pound of body weight diluted with 1 part of the drug to 5 parts of water.
- (3) Federal law restricts this drug to use by or on the order of a licensed veterinarian.
- [44 FR 12993, Mar. 9, 1979, as amended at 50 FR 41489, Oct. 11, 1985]

§ 520.580 Dichlorophene and toluene capsules.

(a) *Specifications*. Each soft gelatin capsule contains 50 milligrams of dichlorophene and 60 milligrams of toluene or multiples thereof. ¹

- (b) *Sponsor*. (1) For single dose only, see 000010, 000061, 000115, 000842, 011615, 017135, 023851, 049968, 050906, and 058670 in §510.600(c) of this chapter.
- (2) For single and multiple dose, see 000010, 000061, and 038782 in 510.600(c) of this chapter.
- (c) Required statement. Consult your veterinarian for assistance in the diagnosis, treatment, and control of parasitism, and before administering to weak or debilitated animals.
- (d) Conditions of use—(1) Amount. (i) Single dose of 100 milligrams of dicholorophene and 120 milligrams of toluene per pound of body weight.¹
- (ii) Divided dose of 100 milligrams of dichlorophene and 120 milligrams of toluene per 5 pounds of body weight (20 and 24 milligrams per pound) daily for 6 days.¹
- (2) Indications for use. It is used for the removal of ascarids (Toxocara canis and Toxascaris leonina) and hookworms (Ancylostoma caninum and Uncinaria stenocephala) and as an aid in removing tapeworms (Taenia pisiformis, Dipylidium caninum, and Echinococcus granulosus) from dogs and cats. 1
- (3) Limitations. Withhold solid foods and milk for at least 12 hours prior to medication and for 4 hours afterward. Repeat treatment in 2 to 4 weeks in animals subject to reinfection.¹

[45 FR 10332, Feb. 15, 1980]

EDITORIAL NOTE: For FEDERAL REGISTER citations affecting §520.580, see the List of CFR Sections Affected, which appears in the Finding Aids section of the printed volume and on GPO Access.

$\S 520.581$ Dichlorophene tablets.

- (a) Specifications. Each tablet contains 1 gram of dichlorophene.
- (b) *Sponsor*. See 023851 in §510.600(c) of this chapter.
- (c) Required statement. Consult your veterinarian for assistance in the diagnosis, treatment, and control of parasitism, and before administering to weak or debilitated animals.
- (d) Conditions of use. Dogs—(1) Amount. Single dose of 1 tablet (1 gram of dichlorophene) for each 10 pounds of body weight.

¹These conditions are NAS/NRC reviewed and deemed effective. Applications for these uses need not include effectiveness data as

specified by \$514.111 of this chapter, but may require bioequivalency and safety information